

THE LANCET

HIV

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Wohl DA, Yazdanpanah Y, Baumgarten A, et al. Bictegravir combined with emtricitabine and tenofovir alafenamide versus dolutegravir, abacavir, and lamivudine for initial treatment of HIV-1 infection: week 96 results from a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet HIV* 2019; published online May 5. [http://dx.doi.org/10.1016/S2352-3018\(19\)30077-3](http://dx.doi.org/10.1016/S2352-3018(19)30077-3).

Appendices

We thank the following GS-US-380-1489 Investigators and their staff: Albrecht H (Palmetto Health Richland, SC, US), Angel J (The Ottawa Hospital - General Campus, CN), Antela Lopez A (Hospital Clinico Universitario de Santiago, ES), Antinori A (IRCCS Istituto Nazionale per le Malattie Infettive "Lazzaro Spallanzani", IT), Arastéh K (EPIMED GmbH, DE), Arribas Lopez JR (Hospital Universitario La Paz, ES), Asmuth DM (CARES Community Health, CA, US), Bartczak J (Rowan Tree Medical, P.A., FL, US), Baumgarten A (Zibp Zentrum für Infektiologie Berlin Prenzlauer Berg GmbH, DE), Benson P (Be Well Medical Center, MI, US), Berhe M (North Texas Infectious Diseases Consultants, PA, TX, US), Bordon J (Providence Hospital – DC, Washington DC, US), Brar I (Henry Ford Hospital, MI, US), Brinson C (Central Texas Clinical Research, TX, US), Brunetta J (Maple Leaf Research/Maple Leaf Medical Clinic, CN), Charest L (Clinique Medicale L'Actuel, CN), Cindrich R (Bronx Lebanon Hospital Center, NY, US), Clarke Ae (Brighton and Sussex University Hospitals NHS Trust, UK), Clotet Sala B (Hospital Universitari Germans Trias i Pujol, ES), Cook PP (East Carolina University, The Brody School of Medicine, Division of Infectious Diseases, NC, US), Cotte L (Hôpital de la Croix Rousse, France), Coulston DR (Premier Clinical Research, WA, US), Crofoot Jr GE (Gordon E. Crofoot MD PA, TX, US), Cruickshank FA (Rosedale Infectious Diseases, NC, US), Cunningham DL (Pueblo Family Physicians, AZ, US), Daar E (Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, CA, US), De Wet JJ (Spectrum Health, CN), DeJesus E (Orlando Immunology Center, FL, US), Dietz C (Kansas City CARE Clinic, MO, US), Edelstein H (Highland Hospital - Alameda Health System, CA, US), Estrada Perez V (Hospital Clinico San Carlos, ES), Fichtenbaum C (University of Cincinnati College of Medicine, Clinical Trials Unit, Division of Infectious Diseases, OH, US), Flamm J (Kaiser Permanente, CA, US), Gallant JE (Southwest CARE Center, NM, US), Gathe Jr, JC (Therapeutic Concepts, TX, US), Girard P-M (Hopital Saint-Antoine, France), Grossberg R (Montefiore Medical Center, NY, US), Gupta SK (Indiana University, Infectious Diseases Research, IN, US), Hagins DP (Chatham County Health Department, GA, US), Hardy WD (Whitman-Walker Health, Washington, DC, US), Hare CB (Kaiser Permanente Medical Center, Clinical Trials Unit, CA, US), Henry WK (Hennepin County

Medical Center, Positive Care Clinic, MN, US), Hileman C (MetroHealth Medical Center, OH, US), Hite AA (Summa Health System, C.A.R.E. Center, OH, US), Hsiao C-B (Positive Health Clinic - Federal North Building, PN, US), Jain M (UT Southwestern Medical Center, Amelia Court HIV Research Clinic, TX, US), Johnson MA (Royal Free London NHS Foundation Trust, UK), Kasper K (Winnipeg Regional Health Authority - Health Sciences Centre Winnipeg, CN), Kinder CA (AHF-Kinder Medical Group, FL, US), Klein D (Kaiser Permanente, Department of Infectious Diseases, CA, US), Koenig E (Instituto Dominicano de Estudios Virologicos (IDEV), Zona Universitaria, DO), LaMarca A (Therafirst Medical Center, FL, US), Lazzarin A (IRCCS Ospedale San Raffaele - Dipartimento di Malattie Infettive, IT), LeBlanc RP (Clinique OPUS Inc., CN), Lebouché B (Chronic Viral Illness Service/McGill University Health Centre (MUHC), CN), Lucasti C (South Jersey Infectious Disease, NJ, US), Maggiolo F (ASST Papa Giovanni XXII, Piazza OMS 1, IT), Mallolas Masferrer J (Hospital Clinic de Barcelona, ES), Mayer CA (St. Joseph's Comprehensive, Research Institute, FL, US), McDonald C (Tarrant County Infectious Disease Associates, TX, US), McGowan J (North Shore University Hospital/Division of Infectious Diseases, NY, US), Meybeck A (CH de Tourcoing, France), Mills A (Mills Clinical Research, CA, US), Miralles Alvarez C (Hospital Alvaro Cunqueiro, ES), Molina J-M (Hopital Saint-Louis, France), Moreno Guillen S (Hospital Universitarion Ramon y Cajal, ES), Mounzer K (Philadelphia FIGHT, PN, US), Newman CL (Augusta University, GA, US), Orkin C (Barts Health NHS Trust, Ambrose King Centre, Royal London Hospital, UK), Osiyemi O (Triple O Research Institute, P.A., FL, US), Palmieri PJ (Upstate Infectious Diseases Associates, LLP, NY, US), Para M (The Ohio State University Medical Center, OH, US), Parks DA (Southampton Clinical Research Group, INC, MO, US), Parsons C (University Medical Center - New Orleans, LA, US), Petroll AE (Medical College of Wisconsin, WI, US), Pierone Jr G (AIDS Research and Treatment Center of the Treasure Coast, FL, US), Podzamczar D (Hospital Universitari de Bellvitge, ES), Polk C (ID Consultants, NC, US), Portilla Sogorb J (Hospital General Universitario de Alicante, ES), Post FA (King's College Hospital, NHS Foundation Trust, UK), Pozniak AL (Chelsea & Westminster NHS Trust, UK), Prelutsky DJ (Southampton Healthcare, Inc, MO, US), Pugliese P (CHU Nice - Hopital Archet 1, France), Rachlis A (Sunnybrook Health

Sciences Centre , CN), Ramgopal M (Midway Immunology and Research Center, FL, US), Rashbaum B (Capital Medical Associates. PC, Washington, DC, US), Richmond GJ (Gary J. Richmond, M.D., P.A., FL, US), Rivero Roman A (Hospital Universitario Reina Sofia, ES), Roberts A (Medical Faculty Associates , Washington, DC, US), Rockstroh J (Universitätsklinikum Bonn (AoR), DE), Ross JDC (Queen Elizabeth Medical Centre, Edgbaston, UK), Ruane PJ (Ruane Clinical Research Group Inc., CA, US), Rubio R (Hospital Universitario 12 de Octubre, ES), Saag MS (University of Alabama Birmingham, AL, US), Salazar L (St. Jude Hospital Yorba Linda DBA, St. Joseph Heritage Healthcare, CA, US), Santana-Bagur JL (Proyecto ACTU, School of Medicine, University of Puerto Rico, PR, US), Santiago Colon L (HOPE Clinical Research, PR, US), Schrader S (Research Access Network, TX, US), Scribner A (Clinic of Longview - Center for Clinical Research, TX, US), Shalit P (Peter Shalit, M.D., WA, US), Shamblaw D (La Playa Medical Group and Clinical Research, CA, US), Shon ASY (Erie County Medical Center, University at Buffalo, NY, US), Sims III J (St. Hope Foundation, TX, US), Slim J (Saint Michael's Medical Center, NJ, US), Stellbrink H-J (ICH Study Center, DE), Stephan C (Universitätsklinikum Frankfurt, Medizinische Klinik II Schwerpunkt Infektiologie, DE), Tebas-Medrano P (Perelman Center for Advanced Medicine at the Hospital of the University of Pennsylvania, PN, US), Thompson MA (AIDS Research Consortium of Atlanta, GA, US), Towner WJ (Kaiser Permanente Los Angeles Medical Center, Infectious Disease Department, CA, US), Ustianowski AP (Pennine Acute Hospitals NHS Trust, Royal Manchester General Hospital, UK), Van Dam C (Cone Health/Regional Center for Infectious Disease, NC, US), Vandekerckhove L (UZ Gent, Belgium), Vandercam B (Cliniques Universitaires UCL Saint-Luc, Belgium), Vanig TJ (Spectrum Medical Group, AZ, US), Voskuhl G (AIDS Arms Inc / Trinity Health and Wellness Center, TX, US), Wade BH (AHF-Pensacola, FL, US), Walker DK (Baystate Infectious Diseases Clinical Research, MA, US), Waters LJ (Central and North West London, NHS Foundation Trust, UK), Wilkin AM (Wake Forest Baptist Health, NC, US), Wohl D (NC TraCS Institute – CTRC, University of North Carolina at Chapel Hill, NC, US), Wohlfeiler MB (AIDS Healthcare Foundation - South Beach, FL, US), Workowski K (Emory Hospital Midtown Infectious Disease Clinic, GA, US), , Wurapa AK (Infectious Disease

Specialists of Atlanta, GA, US), Yazdanpanah Y (CHU Bichat, France), Young B (Apex Research LLC, CO, US), Zorrilla C (Maternal Infant Studies Center (CEMI), University of Puerto Rico, P, US), Zurawski C (Atlanta ID Group, PC, GA, US).

Appendix Table 1. Virologic outcomes at week 96 in participants ≥50 years of age

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=40)	Dolutegravir, abacavir, and lamivudine (n=41)	Bictegravir, emtricitabine, and tenofovir alafenamide vs Dolutegravir, abacavir, and lamivudine
			Difference in Percentages (CI)[†]
HIV-1 RNA <50 copies per mL	40 (100.0%)	35 (85.4%)	14.1% (0.1% to 28.1%)
HIV-1 RNA ≥50 copies per mL	0	1 (2.4%)	
HIV-1 RNA ≥50 copies per mL in Week 96 window	0	0	
Discontinued Due to Lack of Efficacy	0	0	
Discontinued Due to Other Reasons* and Last Available HIV-1 RNA ≥50 copies per mL	0	1 (2.4%)	
No Virologic Data in Week 96 window	0	5 (12.2%)	
Discontinued Due to AE/Death	0	2 (4.9%)	
Discontinued Due to Other Reasons* and Last Available HIV-1 RNA <50 copies per mL	0	3 (7.3%)	
Missing Data but on Study Drug	0	0	

Data are n (%).

The Week 96 window is between Days 631 and 714 (inclusive).

*Other reasons include subjects who discontinued study drug due to investigator's discretion, subject decision, lost to follow-up, noncompliance with study drug, protocol violation, pregnancy, and study terminated by sponsor.

[†] The difference in percentages of subjects with HIV-1 RNA < 50 copies/mL between treatment groups and its 95% CI were calculated based on the Mantel-Haenszel proportions adjusted by baseline HIV-1 RNA stratum and region stratum.

Appendix Table 2. Virologic outcomes at week 96 in participants with cumulative adherence by pill count <95% at week 96.

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=96)	Dolutegravir, abacavir, and lamivudine (n=120)	Bictegravir, emtricitabine, and tenofovir alafenamide vs Dolutegravir, abacavir, and lamivudine
			Difference in Percentages (CI)[†]
HIV-1 RNA <50 copies per mL	71 (74.0%)	103 (85.8%)	-11.1% (-22.1% to 0.0%)
HIV-1 RNA ≥50 copies per mL	0	5 (4.2%)	
HIV-1 RNA ≥50 copies per mL in Week 96 window	0	2 (1.7%)	
Discontinued Due to Lack of Efficacy	0	0	
Discontinued Due to Other Reasons* and Last Available HIV-1 RNA ≥50 copies per mL	0	3 (2.5%)	
No Virologic Data in Week 96 window	25 (26.0%)	12 (10.0%)	
Discontinued Due to AE/Death	1 (1.0%)	2 (1.7%)	
Discontinued Due to Other Reasons* and Last Available HIV-1 RNA <50 copies per mL	22 (22.9%)	7 (5.8%)	
Missing Data but on Study Drug	2 (2.1%)	3 (2.5%)	

Data are n (%).

The Week 96 window is between Days 631 and 714 (inclusive).

*Other reasons include subjects who discontinued study drug due to investigator's discretion, subject decision, lost to follow-up, noncompliance with study drug, protocol violation, pregnancy, and study terminated by sponsor.

† The difference in percentages of subjects with HIV-1 RNA < 50 copies/mL between treatment groups and its 95% CI were calculated based on the Mantel-Haenszel proportions adjusted by baseline HIV-1 RNA stratum and region stratum.

Appendix Table 3. Treatment differences in HIV-1 RNA <50 copies per mL at week 96 by subgroup

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)	Dolutegravir, abacavir, and lamivudine (n=315)	Test for interaction P-Value[†]
Overall	276 (87.9%)	283 (89.8%)	
Age (Years)			N/A [‡]
< 50	236/274 (86.1%)	248/274 (90.5%)	
≥ 50	40/40 (100.0%)	35/41 (85.4%)	
Sex			0.76
Male	251/285 (88.1%)	255/282 (90.4%)	
Female	25/29 (86.2%)	28/33 (84.8%)	
Race			0.91
Black	98/114 (86.0%)	99/112 (88.4%)	
Nonblack	176/198 (88.9%)	184/203 (90.6%)	
Baseline HIV-1 RNA (copies/mL)			0.98
≤ 100,000	233/261 (89.3%)	241/265 (90.9%)	
> 100,000	43/53 (81.1%)	42/50 (84.0%)	
Baseline CD4 Cell Count (/uL)			0.97
< 200	28/36 (77.8%)	26/32 (81.3%)	
≥ 200	248/278 (89.2%)	257/283 (90.8%)	
Region			0.76
US	200/228 (87.7%)	208/233 (89.3%)	
Ex-US	76/86 (88.4%)	75/82 (91.5%)	
Study Drug Adherence (%)			0.029
< 95	71/96 (74.0%)	103/120 (85.8%)	
≥ 95	205/216 (94.9%)	180/195 (92.3%)	

The Week 96 window is between Days 631 and 714 (inclusive).

[†]Test for interaction based on the logistic regression model included baseline HIV-1 RNA stratum (≤ 100,000 vs. > 100,000 copies/mL) and region stratum (US vs. Ex-US) (if not the subgroup factor), subgroup, treatment, and the interaction between treatment and subgroup. P-value for the homogeneity test was from the Wald test of the interaction between treatment and subgroup.

[‡]P-value could not be calculated for age subgroup.

Appendix Table 4. Study drug-related adverse events in ≥1 participant in either group

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)	Dolutegravir, abacavir, and lamivudine (n=315)
Any study drug-related adverse event	89 (28.3%)	127 (40.3%)
Nausea	18 (5.7%)	55 (17.5%)
Diarrhoea	19 (6.1%)	13 (4.1%)
Headache	16 (5.1%)	16 (5.1%)
Fatigue	9 (2.9%)	11 (3.5%)
Abnormal dreams	8 (2.5%)	8 (2.5%)
Dizziness	7 (2.2%)	9 (2.9%)
Insomnia	6 (1.9%)	9 (2.9%)
Abdominal distension	5 (1.6%)	5 (1.6%)
Abdominal pain	3 (1.0%)	6 (1.9%)
Vomiting	3 (1.0%)	6 (1.9%)
Abdominal discomfort	3 (1.0%)	4 (1.3%)
Constipation	4 (1.3%)	2 (0.6%)
Sleep disorder	1 (0.3%)	5 (1.6%)
Dyspepsia	1 (0.3%)	4 (1.3%)
Flatulence	3 (1.0%)	2 (0.6%)
Nightmare	3 (1.0%)	2 (0.6%)
Somnolence	2 (0.6%)	3 (1.0%)
Anxiety	1 (0.3%)	3 (1.0%)
Decreased appetite	1 (0.3%)	3 (1.0%)
Faeces soft	2 (0.6%)	1 (0.3%)
Hot flush	2 (0.6%)	1 (0.3%)
Hyperhidrosis	1 (0.3%)	2 (0.6%)
Myalgia	1 (0.3%)	2 (0.6%)
Neutropenia	2 (0.6%)	1 (0.3%)
Weight increased	0	3 (1.0%)
Alanine aminotransferase increased	2 (0.6%)	0
Aspartate aminotransferase increased	2 (0.6%)	0
Depressed mood	2 (0.6%)	0
Depression	0	2 (0.6%)
Flushing	0	2 (0.6%)
Frequent bowel movements	0	2 (0.6%)
Gamma-glutamyltransferase increased	2 (0.6%)	0
Gastrooesophageal reflux disease	0	2 (0.6%)
Hyperlipidaemia	1 (1 (0.3%)
Low density lipoprotein increased	2 (0.6%)	0
Night sweats	2 (0.6%)	0
Palpitations	2 (0.6%)	0

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)	Dolutegravir, abacavir, and lamivudine (n=315)
Pollakiuria	2 (0.6%)	0
Rash	1 (0.3%)	1 (0.3%)
Rash generalised	1 (0.3%)	1 (0.3%)
Abortion spontaneous	1 (0.3%)	0
Aerophagia	1 (0.3%)	0
Ageusia	0	1 (0.3%)
Arthralgia	1 (0.3%)	0
Asthenia	0	1 (0.3%)
Blood alkaline phosphatase increased	1 (0.3%)	0
Blood cholesterol increased	1 (0.3%)	0
Blood creatine phosphokinase increased	0	1 (0.3%)
Blood creatinine increased	0	1 (0.3%)
Blood lactate dehydrogenase increased	0	1 (0.3%)
Blood uric acid increased	1 (0.3%)	0
Confusional state	0	1 (0.3%)
Cough	0	1 (0.3%)
Creatinine renal clearance decreased	1 (0.3%)	0
Dermatitis	0	1 (0.3%)
Dermatitis contact	0	1 (0.3%)
Disorientation	1 (0.3%)	0
Disturbance in attention	0	1 (0.3%)
Dizziness postural	1 (0.3%)	0
Drug eruption	0	1 (0.3%)
Dry mouth	0	1 (0.3%)
Dysaesthesia	0	1 (0.3%)
Dysgeusia	1 (0.3%)	0
Epigastric discomfort	0	1 (0.3%)
Feeling hot	0	1 (0.3%)
Feeling jittery	0	1 (0.3%)
Folliculitis	1 (0.3%)	0
Food allergy	1 (0.3%)	0
Gastroenteritis	0	1 (0.3%)
Generalised tonic-clonic seizure	1 (0.3%)	0
Hyperbilirubinaemia	1 (0.3%)	0
Hypercholesterolaemia	1 (0.3%)	0
Hypoaesthesia	0	1 (0.3%)
Increased appetite	0	1 (0.3%)
Joint stiffness	0	1 (0.3%)
Keratoconus	0	1 (0.3%)
Libido decreased	1 (0.3%)	0

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)	Dolutegravir, abacavir, and lamivudine (n=315)
Lipodystrophy acquired	0	1 (0.3%)
Lumbar radiculopathy	0	1 (0.3%)
Malaise	0	1 (0.3%)
Middle insomnia	1 (0.3%)	0
Nail bed disorder	1 (0.3%)	0
Osteoporosis	0	1 (0.3%)
Pancreatitis acute	0	1 (0.3%)
Paraesthesia	0	1 (0.3%)
Parkinson's disease	1 (0.3%)	0
Pruritus generalised	1 (0.3%)	0
Psychomotor hyperactivity	0	1 (0.3%)
Retching	0	1 (0.3%)
Rhinorrhoea	1 (0.3%)	0
Skin mass	1 (0.3%)	0
Steatorrhoea	0	1 (0.3%)
Sudden death	1 (0.3%)	0
Tension headache	0	1 (0.3%)
Thirst	1 (0.3%)	0
Thrombocytopenia	0	1 (0.3%)
Tremor	0	1 (0.3%)
Vertigo		1 (0.3%)

Data are n (%). Relatedness to study drug assessed by the investigator.

Appendix Table 5. Grade 3 or 4 laboratory abnormalities in $\geq 2\%$ of participants in either group through week 96

	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)	Dolutegravir, abacavir, and lamivudine (n=315)
Any Grade 3 or 4 Treatment-Emergent Toxicity Grade	71/314 (22.6%)	62/315 (19.7%)
Neutrophils (Decreased)	9/314 (2.9%)	12/315 (3.8%)
Amylase (Increased)	8/314 (2.5%)	10/315 (3.2%)
AST (Increased)	13/314 (4.1%)	8/315 (2.5%)
Creatine Kinase (Increased)	20/314 (6.4%)	16/315 (5.1%)
LDL (Fasting, Increased)	10/308 (3.2%)	11/310 (3.5%)


Appendix Table 6. Changes from Baseline in Fasting Metabolic Laboratory Parameters at Week 96 (Safety Analysis Set)

Metabolic Assessment	Bictegravir, emtricitabine, and tenofovir alafenamide (n=314)		Dolutegravir, abacavir, and lamivudine (n=315)		p-value
	n	Median (Q1, Q3)	n	Median (Q1, Q3)	
Total cholesterol (mg/dL)					
Baseline	305	159 (133, 181)	305	162 (138, 186)	0.23
Change at Week 96	271	15 (1, 34)	273	8 (-7, 26)	0.002
Direct LDL (mg/dL)					
Baseline	305	101 (83, 123)	305	101 (84, 126)	0.46
Change at Week 96	270	17 (2, 32)	272	7 (-5, 24)	<0.001
Triglycerides (mg/dL)					
Baseline	305	93 (67, 132)	305	96 (66, 138)	0.74
Change at Week 96	270	8 (-16, 38)	273	6 (-21, 30)	0.28
HDL (mg/dL)					
Baseline	305	42 (34, 51)	305	42 (35, 51)	0.62
Change at Week 96	271	4 (-1, 11)	273	5 (0, 12)	0.17
Total cholesterol to HDL ratio					
Baseline	305	3.7 (3.0, 4.7)	305	3.7 (3.0, 4.6)	0.93
Change at Week 96	271	-0.1 (-0.5, 0.5)	273	-0.2 (-0.7, 0.3)	0.003
Glucose (mg/dL)					
Baseline	310	88 (82, 93)	309	88 (82, 94)	0.68
Change at Week 96	278	4 (-2, 11)	283	4 (-1, 11)	0.63

Only laboratory measurements under fasting status were summarized.

P-values were from the 2-sided Wilcoxon rank sum test to compare the 2 treatment groups.

Appendix Figure 1. Prevalence and incidence of nausea through 96 weeks



B/F/TAF, bicitgravir, emtricitabine, and tenofovir alafenamide; DTG/ABC/3TC, dolutegravir, abacavir, lamivudine